

# Accumulative weekly load in a professional football team: with special reference to match playing time and game position

**AUTHORS:** David Casamichana<sup>1,2</sup>, Andrés Martín-García<sup>3</sup>, Antonio Gómez Díaz<sup>4,5</sup>, Paul S Bradley<sup>6</sup>, Julen Castellano<sup>7</sup>

<sup>1</sup> Atlantic European University. Santander, Spain

<sup>2</sup> Real Sociedad Sports Performance Department, San Sebastian, Spain

<sup>3</sup> FC Barcelona Sports Performance Department. Barcelona, Spain

<sup>4</sup> Murcia University, Faculty of Sports Sciences, San Javier, Spain

<sup>5</sup> Polish Football Federation

<sup>6</sup> Research Institute for Sport & Exercise Sciences, Liverpool John Moores University, Liverpool, UK

<sup>7</sup> University of the Basque Country (UPV/EHU), Spain

**ABSTRACT:** The aim of this study was to compare weekly accumulative load during the in-season competitive period by professional soccer players according to the amount of time played in official matches (90-min, >60-min, <60-min, and 0-min) regarding the players' position. Twenty-four professional outfield football players were monitored by GPS devices during training sessions and official matches and the accumulative weekly load were calculated for the following external load variables: total distance (TD; m), high speed running (HSR; >19.8 km·h<sup>-1</sup>), sprint meters (SPR; >25.2 km·h<sup>-1</sup>), high metabolic load distance (HMLD; >25.5 W·kg<sup>-1</sup>), number of accelerations (ACC; >3 m·s<sup>-2</sup>) and decelerations (DEC; <-3 m·s<sup>-2</sup>). This study revealed that players (as a whole or per demarcation) with more match playing time had a higher accumulative weekly load for most of the variables, but particularly at TD and HMLD (90-min and >60-min vs. <60-min and 0-min), HSR (90-min vs. <60-min and 0-min) and SPR (90-min vs. <60-min and 0-min). In addition, less positional variation was observed in relation to the competition requirements. This information would allow coaches to refine the required load with the objective of optimizing performance to soccer players.

**CITATION:** Casamichana D, Martín-García A, Gómez Díaz A et al. Accumulative weekly load in a professional football team: with special reference to match playing time and game position. *Biol Sport*. 2022;39(1):115-124.

Received: 2020-08-07; Reviewed: 2020-11-07; Re-submitted: 2020-12-11; Accepted: 2021-01-10; Published: 2021-03-04.

Corresponding author:

**Andrés Martín García**

FC Barcelona Sports Performance Department

Av. Onze de Setembre, s/n,

08970 Sant Joan Despí

Barcelona

Ciudad Deportiva Joan Gamper

Tel.: +34 00 660251444

Email: andres.martin@

fcbarcelona.cat

**ORCID:**

David Casamichana

0000-0002-2082-4344

Andrés Martín-García

0000-0002-7341-4664

Antonio Gómez Díaz

0000-0001-7151-6133

Paul S Bradley

Julen Castellano

0000-0001-5167-5284

**Key words:**

Football

GPS

Periodization

Training load

Positional role

## INTRODUCTION

Football (soccer) is a complex sport where a multitude of tactical, technical and physical elements modulate performance [1]. Contemporary approaches examining the integrative nature of the game, in which an array of factors are synthesised, provide greater insights into a player's positional and match requirements [2–3]. These multifaceted demands accumulate across the season, particularly when matches are played with minimal recovery [4]. Accordingly, practitioners modify training and competition demands during the season by alternating load and rotating the squad [5]. In order to achieve a balance between acute and chronic loads, the applied scientist must have an in-depth knowledge of competition and training loads within the microcycle [6]. Thus, enabling optimal conditioning for performance but more importantly identifying players that are inadequately loaded and therefore potentially susceptible to injury [7–8]. Although, some have examined the properties of the training microcycle in relation to the days before competition [9–10–11–12], limited research exists regarding the accumu-

lative load of players achieving match time versus those with limited exposure to competition.

Using global positioning system (GPS) technology in both training and official matches is an effective tool for monitoring the accumulative player load. Despite this technology being widely used within football to quantify training demands, its use in official matches is less common [13]. Thus, limited research exists on the comparison of external loads among players within the microcycle based on official match time. Stevens et al. [14] demonstrated that the load of nonstarter sessions was generally lower than regular training, but this study only provided a basic breakdown of load (starters vs. non starters) without differentiating between match exposure times (e.g. 0-min vs. <60-min vs. >60-min vs. 90-min). Competing in weekly matches seems to act as an important physiological, mechanical and neuromuscular stimulus for various fitness parameters [15–16–17]. Thus, creating a discrepancy in total load between those getting regular match time versus those with partial or no

match time [14]. Furthermore, players completing more match minutes perceived training to be harder the following week compared to those with less match time [18]. However, research has yet to establish if this finding is related to a higher training load or accumulated fatigue. Moreover, the tactical role of a player seems to be a key determinant of their match physical performance, so it is imperative that the conditioning stimulus has a positional element to it [19–20]. As it is common for defensive positions to complete more match time and offensive players to be regularly replaced [21], it might be prudent to determine if position specific trends exist regarding accumulative load. This would enable bespoke conditioning practices to be developed that account for the players weekly load and tactical role in the team. Accordingly, this study aimed to compare weekly accumulative load according to the amount of time played in official matches (e.g., 90-min, >60-min, <60-min and 0-min) and in relation to a players' position.

## MATERIALS AND METHODS

### Participants

Twenty-four professional outfield football players participated in this study (age;  $20 \pm 2$  yr, body mass;  $70.2 \pm 6.1$  kg and stature;  $1.78 \pm 0.1$  m). Players belonged to a reserve squad of a Spanish La Liga club that also competed in the UEFA Champions League. Data were collected during the competitive season as part of the players' daily monitoring. Outfield players not completing a full microcycle were excluded from the analysis. Data arose as a condition of the players' employment whereby they were assessed daily, thus no authorization was required from an institutional ethics committee [22]. Nevertheless, this study complied with the Declaration of Helsinki and players provided informed consent before participating.

**TABLE 1.** Number of files for each group and position.

	Game Exposure				Total files
	90-min	> 60-min	< 60-min	0-min	
<b>CD</b>	9	2	5	10	26
<b>FB</b>	17	3	4	10	34
<b>MF</b>	6	1	3	2	12
<b>OMF</b>	7	4	7	12	30
<b>FW</b>	9	4	5	2	20
<b>Total</b>	48	14	24	36	122

CD = central defender; FW = forward; MF = midfielder; WMF = wide midfielder; FB = full back; 0-min: group of players with 0 minutes in competition; < 60-min: group of players with less than 60 minutes in competition; > 60-min: group of players with more than 60 minutes in competition; 90-min: group of players with 90 minutes in competition.

### Design

GPS data were collected from 37 competitive matches and 42 training weeks during the 2015–16 season. This enabled absolute and relative external training loads to be quantified across the microcycle for players accumulating various match exposures. Sessions were performed on a natural grass surface within a pitch dimension of  $105 \times 68$  m. The team systematically played in a 4–3–3 formation, with two full backs (FB), two central defenders (CD), one midfielder (MD), two offensive midfielders (OMF) and three forwards (FW). The players were grouped by the number of minutes they played in the previous match (Table 1): the full match (90-min), more than 60-min of the match but did not complete it (>60-min), less than 60-min of the match (<60-min) and did not play in the match (0-min).

### Structured Microcycle

Due to variations in the number of days between matches [12–14], the present study only analysed training weeks where players had six days between successive matches and the training week composed of five training sessions that had a clear focus on an upcoming match [23]. Based on the recommendations of Akenhead et al. [13], training load data were qualified with respect to the number of days before or after a match (MD minus or plus). The training sessions that are contextualised below were composed of integrated content (e.g., tactical, technical and physical factors were amalgamated):

*MD+1*: was the session the day after competition where players were split into two training groups. The first group included players who had played >60-min of previous match (90-min and >60-min). The aim of this session was to recover after the match, so the term “recovery session” was used, MD+1 recovery (MD+1R). These players conducted low-impact exercises combined with regeneration exercises. The second group included players who had played <60-min of previous match (<60-min and 0-min). This group worked within a technical circuit followed by a small positional game and a small-sided game (SSG) with goalkeepers within an assigned area which ranged between 30 and 60 m<sup>2</sup> per player. This session attempted to reproduce competitive loads, so the term “compensatory session” was used, MD+1 compensatory (MD+1C). The session MD-4, MD-3, MD-2 and MD-1 were common for players, regardless of the number of minutes played in previous official match. The content of each of the sessions has been described by Martin-Garcia et al. [11].

### Physical variables

Activity profiles of players were monitored during each match and training session using a portable 10 Hz GPS unit (Viper Pod, 50 g,  $88 \times 33$  mm, Statsports Viper, Northern Ireland). Each unit was placed in a specially designed vest, inside a mini pocket positioned between the shoulder blades. Quantifying the devices accuracy indicated a 2.5% estimation error in distance covered, with accuracy improving as the distance covered increased and the speed of movement decreased [24]. In order to avoid inter-unit error, each player used the same device during the study period [25–26]. Upon com-

## Weekly external load in soccer

**TABLE 2.** Absolute accumulated values per external load variables (mean ± SD) obtained from each player's group during a training microcycle plus match.

Variables	Position	0-min	< 60-min	> 60-min	90-min	ES;p
TD	CD	21957.6 ± 1764.8	25945.6 ± 3179.9	29387.9 ± 5027.4	30378.1 ± 1323.3*	ES: 0.4–5.4; p < 0.001
	FB	22657.5 ± 3051.8	27200.0 ± 2714.8*	28906.7 ± 1855.4*	32093.4 ± 1998.5 <sup>e,*†</sup>	ES: 1.6–3.8; p < 0.001
	MF	23423.1 ± 1603.9	25643.6 ± 5250.0	33700.2 ± 2603.2 <sup>e,*†</sup>	32143.1 ± 1474.0 <sup>e,*</sup>	ES: 0.8–4.9; p = 0.002
	OMF	23604.9 ± 2896.2	24064.6 ± 5164.0	32296.7 ± 3410.1 <sup>e,*†</sup>	33930.5 ± 1672.6 <sup>e,*†</sup>	ES: 0.7–4.1; p < 0.001
	FW	19983.4 ± 2519.1	24670.5 ± 3436.0*	27395.2 ± 728.0*	28006.1 ± 2967.1	ES: 0.2–2.8; p = 0.012
	AVG	22715.6 ± 2644.8	25328.4 ± 3899.6*	29690.6 ± 2755.3 <sup>*†</sup>	31327.2 ± 2656.9 <sup>*†</sup>	ES: 0.6–3.2; p < 0.001
ES;p	ES: 0.1–1.3; p = 0.335	ES: 0.4–0.7; p = 0.773	ES: 0.4–4.1; p = 0.101	ES: 1.1–2.4; p < 0.001	.	
HSR	CD	537.2 ± 119.5	806.1 ± 295.2	821.1 ± 16.8*	1027.4 ± 245.1*	ES: 0.9–2.5; p < 0.001
	FB	991.0 ± 456.9 <sup>c</sup>	1479.1 ± 240.1 <sup>a,c,d</sup>	1477.4 ± 462.1 <sup>a,e</sup>	1625.4 ± 384.0 <sup>a,c,e,*</sup>	ES: 0.4–1.5; p = 0.003
	MF	434.4 ± 124.6	582.5 ± 180.3	1152.9 ± 266.6 <sup>*†</sup>	837.2 ± 230.5*	ES: 1.3–3.7; p = 0.027
	OMF	676.6 ± 263.9	636.1 ± 261.0	1284.6 ± 298.8 <sup>*†</sup>	1289.6 ± 388.2 <sup>*†</sup>	ES: 0.0–1.9; p < 0.001
	FW	466.2 ± 98.2	863.0 ± 452.4	969.0 ± 177.1	1221.3 ± 314.0*	ES: 0.9–2.6; p = 0.037
	AVG	700.1 ± 348.5	850.7 ± 409.8	1147.4 ± 244.5*	1277.9 ± 427.9 <sup>*†</sup>	ES: 0.3–1.5; p < 0.001
ES;p	ES: 0.9–1.4; p = 0.007	ES: 1.6–4.1; p = 0.003	ES: 0.5–1.8; p = 0.124	ES: 0.9–2.3; p < 0.001	.	
SPR	CD	104.8 ± 65.9	119.1 ± 93.4	134.6 ± 22.8	202.0 ± 95.0	ES: 0.8–1.2; p = 0.068
	FB	225.5 ± 124.1 <sup>c,d,e</sup>	374.5 ± 116.8 <sup>d</sup>	324.1 ± 160.6 <sup>a,c,d,e</sup>	381.3 ± 147.4 <sup>a,c,d,*</sup>	ES: 0.1–1.1; p = 0.043
	MF	57.6 ± 29.7	62.9 ± 24.8	184.8 ± 30.7*	77.6 ± 34.9	ES: 3.1–4.2; p = 0.034
	OMF	86.3 ± 61.5	62.2 ± 61.5	139.1 ± 73.4	200.9 ± 80.4 <sup>*†</sup>	ES: 0.8–1.9; p = 0.002
	FW	35.9 ± 13.3	193.4 ± 174.9	170.0 ± 119.8	247.6 ± 113.9 <sup>c</sup>	ES: 0.4–2.1; p = 0.292
	AVG	125.5 ± 103.2	152.2 ± 147.8	188.9 ± 81.5	254.0 ± 150.5 <sup>*†</sup>	ES: 0.5–1.0; p < 0.001
ES;p	ES: 1.2–1.7; p = 0.001	ES: 1.2–3.6; p = 0.002	ES: 1.1–1.5; p = 0.290	ES: 1.0–2.4; p < 0.001	.	

TABLE 2. Continue.

Variables	Position	0-min	< 60-min	> 60-min	90-min	ES;p
	CD	2716.9 ± 229.0	3454.1 ± 513.5*	3684.4 ± 315.5*	3891.7 ± 325.2*	ES: 0.6–4.1; p < 0.001
	FB	3381.5 ± 948.4 <sup>e</sup>	4389.8 ± 403.4	4310.4 ± 668.5	4958.2 ± 702.3 <sup>a,c,e,*</sup>	ES: 0.9–2.0; p < 0.001
	MF	3075.9 ± 114.4	3553.7 ± 908.9	5077.1 ± 304.4 <sup>a,e,*†</sup>	4023.7 ± 547.3 <sup>*†</sup>	ES: 2.0–9.4; p = 0.054
<b>HMLD</b>	OMF	3233.9 ± 614.2 <sup>e</sup>	3224.3 ± 1032.0	5315.5 ± 861.6 <sup>a,b,e,*†</sup>	5125.7 ± 707.1 <sup>a,c,e,*†</sup>	ES: 0.3–2.5; p < 0.001
	FW	2506.3 ± 133.3	3413.4 ± 506.4	3882.5 ± 245.0*	3940.2 ± 683.1*	ES: 0.1–2.3; p = 0.019
	AVG	3093.5 ± 675.0	3543.3 ± 779.6	4403.4 ± 479.1 <sup>*†</sup>	4465.8 ± 806.2 <sup>*†</sup>	ES: 0.1–1.8; p < 0.001
	<b>ES;p</b>	<b>ES: 0.2–1.0; p = 0.119</b>	<b>ES: 1.2–1.4; p = 0.188</b>	<b>ES: 0.3–2.2; p = 0.025</b>	<b>ES: 0.2–1.7; p &lt; 0.001</b>	.
	CD	530.8 ± 71.8	633.6 ± 84.5*	586.5 ± 118.0	621.2 ± 37.7*	ES: 0.2–1.3; p = 0.014
	FB	613.0 ± 152.4	721.0 ± 19.2	652.3 ± 130.8	676.1 ± 155.4	ES: 0.3–0.8; p = 0.624
	MF	626.6 ± 28.3	705.3 ± 162.6	796.0 ± 108.3 <sup>e</sup>	630.9 ± 101.6	ES: 0.6–2.4; p = 0.445
<b>ACC</b>	OMF	521.9 ± 160.4	552.4 ± 192.1	783.5 ± 143.2 <sup>e,*</sup>	737.9 ± 100.2 <sup>e,*</sup>	ES: 0.4–1.7; p = 0.005
	FW	405.0 ± 82.0	452.0 ± 92.3	491.6 ± 54.2	514.3 ± 133.0	ES: 0.2–0.9; p = 0.526
	AVG	551.9 ± 137.0	597.2 ± 154.0	634.5 ± 110.9	640.6 ± 142.6*	ES: 0.0–0.6; p = 0.020
	<b>ES;p</b>	<b>ES: 0.1–3.9; p = 0.180</b>	<b>ES: 0.1–3.8; p = 0.033</b>	<b>ES: 0.1–4.1; p = 0.023</b>	<b>ES: 0.4–1.9; p = 0.011</b>	.
	CD	501.4 ± 57.6	574.6 ± 80.4	498.0 ± 131.5	553.7 ± 42.3	ES: 0.4–0.8; p = 0.098
	FB	588.4 ± 145.7	652.5 ± 12.7 <sup>e</sup>	637.7 ± 132.3 <sup>a,e</sup>	688.9 ± 134.5 <sup>a,e</sup>	ES: 0.3–0.7; p = 0.281
	MF	629.0 ± 30.3 <sup>a,e</sup>	659.3 ± 155.9	767.0 ± 120.7 <sup>a,e</sup>	593.6 ± 63.1	ES: 0.8–2.2; p = 0.307
<b>DEC</b>	OMF	500.0 ± 153.4	486.0 ± 165.9	767.0 ± 115.2 <sup>a,e,*†</sup>	680.4 ± 73.0 <sup>a,e,*</sup>	ES: 1.0–1.9; p = 0.002
	FW	442.0 ± 90.5	478.6 ± 69.9	464.6 ± 38.1	507.0 ± 92.9	ES: 0.3–0.7; p = 0.590
	AVG	532.2 ± 129.4	553.3 ± 129.3	604.5 ± 107.5	616.1 ± 119.1*	ES: 0.1–0.7; p = 0.008
	<b>ES;p</b>	<b>ES: 0.3–3.0; p = 0.191</b>	<b>ES: 0.1–1.6; p = 0.071</b>	<b>ES: 0.0–3.7; p = 0.007</b>	<b>ES: 0.1–1.2; p &lt; 0.001</b>	.

CD = central defender; FW = forward; MF = midfielder; WMF = wide midfielder; FB = full back; 0-min: group of players with 0 minutes in competition; < 60-min: group of players with less than 60 minutes in competition; > 60-min: group of players with more than 60 minutes in competition; 90-min: group of players with 90 minutes in competition; TD = distance covered; HSR = high speed running ( $m > 19.8 \text{ km}\cdot\text{h}^{-1}$ ); SPR = sprint ( $m > 25.2 \text{ km}\cdot\text{h}^{-1}$ ); HMLD = high metabolic load distance (m); ACC = accelerations ( $n > 3 \text{ m}\cdot\text{s}^{-2}$ ); DEC = decelerations ( $n < -3 \text{ m}\cdot\text{s}^{-2}$ ).

pletion of each match and session, GPS data were extracted using proprietary software (Viper, Statsports, Ireland). The total distance (TD; m), high-speed running (HSR;  $m > 19.8 \text{ km}\cdot\text{h}^{-1}$ ) and sprinting (SPR;  $m > 25.2 \text{ km}\cdot\text{h}^{-1}$ ) distances were quantified along with the number of intense accelerations/decelerations (ACC/DEC;  $> 3 \text{ m}\cdot\text{s}^{-2}$ ), and high metabolic load distance (HMLD;  $m > 25.5 \text{ W}\cdot\text{kg}^{-1}$ ). The mean value of each training group (90-min;  $> 60$ -min;  $< 60$ -min, and 0-min) was expressed in absolute values and the estimated accumulative load in relation to the mean external load registered during official matches. In the case of players without official match records, the mean values of the position they occupy in the game system were applied (e.g. three offensive midfielders and one central defender).

*Statistical analysis*


All statistical analyses were conducted using SPSS for Windows 16.0 (SPSS Inc., Chicago, USA). Homogeneity of variance was examined by conducting Levene's test. A one-way ANOVA was used to evaluate differences in dependent variables across various groups (90-min,  $> 60$ -min,  $< 60$ -min, and 0-min). This analysis also was implemented regarding player' positions (CD, FW, MF, WMF and FB). In the event of a difference occurring, Bonferroni post-hoc tests were used to identify any localised effects or Dunnett's T3 post-hoc test was applied when the variances were not homogeneous. The data are presented as means and standard deviations and alpha was set at  $p < 0.05$ . Effect sizes (ES) were calculated to determine mean-

ingful differences. Magnitudes of difference were classed as [27]: trivial ( $< 0.2$ ), small ( $> 0.2-0.6$ ), moderate ( $> 0.6-1.2$ ), large ( $> 1.2-2.0$ ), and very large ( $> 2.0-4.0$ ).

**RESULTS**

*Absolute Accumulative Load*

Table 2 presents the absolute values obtained from each training group (90-min,  $> 60$ -min,  $< 60$ -min, and 0-min) across variables. The 0-min group had a lower accumulative load than the 90-min group for TD, HSR, SPR, HMLD, ACC, DEC ( $p < 0.01$ , ES: 0.7-3.2) and the  $> 60$ -min group for TD, HSR, HMLD ( $p < 0.01$ , ES: 1.3-2.5). The  $< 60$ -min group had a lower accumulative load than the 90-min group for TD, HSR, SPR and HMLD ( $p < 0.01$ , ES: 0.7-1.9). The  $> 60$ -min group had a higher accumulative load than  $< 60$ -min group for TD, HMLD ( $p < 0.05$ , ES: 1.1-1.3) and the 0-min group for TD, HSR and HMLD ( $p < 0.01$ , ES: 0.8-2.1). Positional differences were evident across all groups (Table 2), specifically FB in the 0-min group demonstrating consistently higher loads than MF for HSR ( $p < 0.01$ , ES: 1.4) and SPR ( $p < 0.01$ , ES: 1.7) and FW for HMLD ( $p < 0.05$ , ES: 1.0). FB in  $< 60$ -min group also had higher weekly loads than OMF for HSR ( $p < 0.01$ , ES: 3.2) and SPR ( $p < 0.01$ , ES: 3.6). In the  $> 60$ -min group, MF and OMF also demonstrated greater loads than FW for HMLD ( $p < 0.05$ , ES: 0.9-2.2) and ACC ( $p < 0.05$ , ES: 1.0-4.1). For the 90-min group, FB had higher loads than CD and MF for HSR ( $p < 0.01$ , ES: 1.0-2.3) and SPR ( $p < 0.01$ , ES: 1.4-2.4),



**FIG. 1.** Relative accumulative load for each training variable regarding the training groups without considering the position of the players in the system.

0-min: group of players with 0 minutes in competition;  $< 60$ -min: group of players with less than 60 minutes in competition;  $> 60$ -min: group of players with more than 60 minutes in competition; 90-min: group of players with 90 minutes in competition; TD = distance covered; HSR = high speed running ( $m > 19.8 \text{ km}\cdot\text{h}^{-1}$ ); SPR = sprint ( $m > 25.2 \text{ km}\cdot\text{h}^{-1}$ ); HMLD = high metabolic load distance (m); ACC = accelerations ( $n > 3 \text{ m}\cdot\text{s}^{-2}$ ); DEC = decelerations ( $n < -3 \text{ m}\cdot\text{s}^{-2}$ ).

**TABLE 3. Relative accumulated** values per external load variables (mean  $\pm$  SD) for each player's group during a training microcycle plus match.

Variables	Position	0-min	< 60-min	> 60-min	90-min	ES;p
%TD	CD	224.3 $\pm$ 18.5	268.0 $\pm$ 32.9	301.2 $\pm$ 53.6 <sup>e</sup>	312.8 $\pm$ 15.5*	ES: 0.5–5.2; p < 0.001
	FB	228.3 $\pm$ 28.5	279.1 $\pm$ 27.1*	297.0 $\pm$ 19.0*	323.9 $\pm$ 23.1 <sup>c,*†</sup>	ES: 1.2–3.8; p < 0.001
	MF	214.9 $\pm$ 14.7	235.3 $\pm$ 48.2	309.3 $\pm$ 13.4 <sup>e,*</sup>	292.8 $\pm$ 13.5*	ES: 1.2–6.6; p = 0.003
	OMF	210.0 $\pm$ 24.8	225.2 $\pm$ 59.6	299.3 $\pm$ 19.4*	313.4 $\pm$ 22.9 <sup>*†</sup>	ES: 0.6–4.3; p < 0.001
	FW	217.7 $\pm$ 27.4	264.8 $\pm$ 45.7	279.1 $\pm$ 24.9	309.2 $\pm$ 26.7*	ES: 1.2–3.4; p = 0.026
	AVG	219.5 $\pm$ 24.2	253.2 $\pm$ 47.0*	293.0 $\pm$ 26.0 <sup>*†</sup>	313.4 $\pm$ 22.8 <sup>*†</sup>	ES: 0.9–4.0; p < 0.001
	ES;p	ES: 0.2–0.7; p = 0.435	ES: 0.4–1.1; p = 0.294	ES: 0.2–1.4; p = 0.705	ES: 0.4–1.5; p = 0.036	.
%HSR	CD	120.1 $\pm$ 25.9	182.0 $\pm$ 72.1	196.0 $\pm$ 35.3	234.8 $\pm$ 64.9*	ES: 0.6–2.2; p = 0.002
	FB	139.1 $\pm$ 50.9	174.8 $\pm$ 20.5	233.3 $\pm$ 65.5*	237.4 $\pm$ 47.7*	ES: 0.1–2.0; p < 0.001
	MF	96.7 $\pm$ 27.7	129.7 $\pm$ 40.1	256.6 $\pm$ 32.1 <sup>*†</sup>	177.4 $\pm$ 38.8*	ES: 2.1–5.4; p = 0.012
	OMF	97.7 $\pm$ 51.5	105.8 $\pm$ 65.3	196.4 $\pm$ 36.3*	205.8 $\pm$ 60.0 <sup>*†</sup>	ES: 0.2–2.0; p < 0.001
	FW	70.6 $\pm$ 14.8	142.1 $\pm$ 54.9	136.1 $\pm$ 47.0	179.3 $\pm$ 27.8*	ES: 0.9–4.2; p = 0.033
	AVG	113.5 $\pm$ 46.6	145.3 $\pm$ 61.7	187.6 $\pm$ 43.2*	214.3 $\pm$ 54.9 <sup>*†</sup>	ES: 0.5–2.0; p < 0.001
	ES;p	ES: 0.5–1.5; p = 0.125	ES: 0.1–1.1; p = 0.187	ES: 0.4–2.8; p = 0.090	ES: 0.1–1.3; p = 0.014	.
%SPR	CD	98.0 $\pm$ 70.7	106.5 $\pm$ 79.7	144.8 $\pm$ 69.0	189.9 $\pm$ 92.2	ES: 0.5–1.1; p = 0.086
	FB	143.7 $\pm$ 72.3	140.8 $\pm$ 26.9	248.3 $\pm$ 79.6 <sup>e,d</sup>	250.0 $\pm$ 83.3 <sup>c,e,*</sup>	ES: 0.0–1.4; p = 0.003
	MF	153.0 $\pm$ 78.8	167.0 $\pm$ 65.8	490.4 $\pm$ 93.5 <sub>a,b,d,e,*†,□</sub>	146.4 $\pm$ 69.6	ES: 4.4–4.6; p = 0.008
	OMF	83.6 $\pm$ 55.3	72.6 $\pm$ 89.6	133.6 $\pm$ 34.6	213.7 $\pm$ 67.1 <sup>*†</sup>	ES: 1.4–1.8; p < 0.001
	FW	21.1 $\pm$ 7.8	143.2 $\pm$ 72.3	80.8 $\pm$ 63.6	150.6 $\pm$ 60.5	ES: 0.1–2.4; p = 0.064
	AVG	106.6 $\pm$ 70.7	117.1 $\pm$ 76.2	164.2 $\pm$ 68.1	201.6 $\pm$ 85.7 <sup>*†</sup>	ES: 0.4–1.2; p < 0.001
	ES;p	ES: 0.1–2.1; p = 0.057	ES: 0.3–1.1; p = 0.326	ES: 2.8–5.6; p < 0.001	ES: 0.5–1.3; p = 0.011	.

TABLE 3. Continue.

	CD	177.3 ± 14.8	226.5 ± 34.8	241.6 ± 12.7*	254.7 ± 21.2*	ES: 0.7–4.2; p < 0.001
	FB	180.9 ± 35.7 <sup>d</sup>	225.4 ± 22.0	246.0 ± 28.1*	270.4 ± 27.1 <sup>c,*</sup>	ES: 0.9–2.9; p < 0.001
	MF	153.8 ± 5.7	177.8 ± 45.5	254.0 ± 33.6*	221.1 ± 30.0*	ES: 1.1–4.6; p = 0.025
%HMLD	OMF	139.7 ± 36.6	163.3 ± 76.4	267.0 ± 60.7 <sup>a,*†</sup>	245.5 ± 35.0 <sup>*†</sup>	ES: 0.5–2.9; p < 0.001
	FW	156.9 ± 8.3	211.1 ± 53.3	213.5 ± 48.5	245.3 ± 26.7	ES: 0.9–3.6; p = 0.131
	AVG	162.5 ± 34.2	199.7 ± 56.4	240.7 ± 36.7*	252.6 ± 31.1 <sup>*†</sup>	ES: 0.3–2.8; p < 0.001
	ES;p	ES: 0.1–1.1; p = 0.019	ES: 0.0–1.0; p = 0.219	ES: 0.2–1.0; p = 0.576	ES: 0.6–0.9; p = 0.004	.
	CD	296.6 ± 42.4	354.1 ± 44.9	331.2 ± 72.6	349.3 ± 23.8*	ES: 0.2–1.3; p = 0.020
	FB	319.0 ± 55.9 <sup>c</sup>	358.6 ± 12.5	352.3 ± 33.6	396.3 ± 38.2 <sup>c,*</sup>	ES: 1.1–1.7; p = 0.001
	MF	250.8 ± 11.3	282.3 ± 65.1	318.6 ± 40.2*	329.9 ± 50.0*	ES: 0.2–1.9; p = 0.168
%ACC	OMF	261.0 ± 92.8	320.3 ± 148.9	420.8 ± 91.4 <sup>a,c,e,*</sup>	362.4 ± 52.5*	ES: 0.9–1.7; p = 0.034
	FW	326.2 ± 66.0 <sup>c</sup>	353.2 ± 59.0	328.7 ± 57.5	382.9 ± 55.7	ES: 0.5–1.0; p = 0.656
	AVG	288.8 ± 70.0*	336.5 ± 87.1*	357.6 ± 59.1*	370.4 ± 48.0*	ES: 0.2–1.4; p < 0.001
	ES;p	ES: 0.1–1.8; p = 0.227	ES: 0.1–1.8; p = 0.752	ES: 0.9–1.3; p = 0.346	ES: 0.3–1.4; p = 0.006	.
	CD	290.9 ± 34.9	339.3 ± 47.2	299.1 ± 98.4	329.3 ± 22.3	ES: 0.3–1.2; p = 0.078
	FB	304.2 ± 56.9	340.8 ± 37.5	334.7 ± 53.7	368.3 ± 45.2 <sup>c,*</sup>	ES: 0.6–1.3; p = 0.018
	MF	248.6 ± 11.9	260.6 ± 61.6	303.2 ± 45.6	299.6 ± 37.8	ES: 0.1–1.8; p = 0.279
%DEC	OMF	237.8 ± 71.7	273.2 ± 124.4	415.9 ± 75.9 <sup>a,c,e,*†</sup>	339.9 ± 37.2*	ES: 1.4–2.5; p = 0.002
	FW	327.7 ± 67.1	346.5 ± 37.2	303.5 ± 68.3	344.6 ± 47.0	ES: 0.0–0.8; p = 0.635
	AVG	275.2 ± 62.8	313.0 ± 80.0	339.1 ± 68.4	342.8 ± 44.4*	ES: 0.1–1.3; p < 0.001
	ES;p	ES: 0.4–1.3; p = 0.040	ES: 0.1–1.8; p = 0.295	ES: 1.2–1.4; p = 0.231	ES: 0.5–1.6; p = 0.005	.

CD = central defender; FW = forward; MF = midfielder; WMF = wide midfielder; FB = full back; 0-min: group of players with 0 minutes in competition; < 60-min: group of players with less than 60 minutes in competition; > 60-min: group of players with more than 60 minutes in competition; 90-min: group of players with 90 minutes in competition; TD = distance covered; HSR = high speed running ( $m > 19.8 \text{ km}\cdot\text{h}^{-1}$ ); SPR = sprint ( $m > 25.2 \text{ km}\cdot\text{h}^{-1}$ ); HMLD = high metabolic load distance (m); ACC = accelerations ( $n > 3 \text{ m}\cdot\text{s}^{-2}$ ); DEC = decelerations ( $n < -3 \text{ m}\cdot\text{s}^{-2}$ ).

while OMF had accumulated more load than FW for HMLD ( $p < 0.01$ , ES: 1.7), ACC ( $p < 0.01$ , ES: 1.9) and DEC ( $p < 0.01$ , ES: 1.2).

### *Relative Accumulative Load*

Figure 1 displays the relative accumulative load for each training variable regarding the training groups without considering the position of the players in the system. The 0-min group had a lower relative accumulative load than the 90-min group for TD, HSR, SPR, HMLD, ACC, DEC ( $p < 0.01$ , ES: 0.7–3.2) and the >60-min group for TD, HSR, HMLD ( $p < 0.05$ , ES: 1.3–2.5). The 90-min group had a higher accumulative load than the 0-min group for TD, HSR, SPR, HMLD, ACC, DEC ( $p < 0.01$ , ES: 1.3–4.5), in addition to the <60 and >60-min groups for TD, HSR, SPR and HMLD ( $p < 0.05$ , ES: 1.0–3.5).

Table 3 presents the accumulative load for each training group (90-min, >60-min, <60-min, and 0-min) relative to match load and across variables and playing positions. Compared to the absolute accumulated load data, less relative differences were evident across position (Table 3). In the 0-min group, FB demonstrated higher accumulative loads than OMF for HMLD ( $p < 0.05$ , ES: 1.1) while FB and FW loaded more than MF for ACC ( $p > 0.05$ , ES: 1.3–1.8). For the >60-min group, OMF accumulated more weekly load than CD, MF and FW for both ACC ( $p > 0.05$ , ES: 1.0–1.3) and DEC ( $p > 0.05$ , ES: 1.2–1.4). Within the 90-min group, FB were consistently loaded more during the week than MF for TD, SPR, HMLD, ACC and DEC ( $p < 0.05$ , ES: 0.9–1.6).

## **DISCUSSION**

This study findings revealed that players with more match playing time had a higher accumulative weekly load for most of the external load variables, but particularly at HSR and SPR. Moreover, position specific accumulative load trends were evident in players completing more match time, with less positional variation when expressed relative to match requirements.

Accumulative weekly load has previously been examined in English [13], Dutch [15] and Spanish League players [18–28]. This is particularly relevant as cultural and competition demands across leagues could result in distinct loading variations in attempt to optimise performance. Interestingly, when comparing external loads across various European leagues, starters consistently display higher accumulative demands than non-starters [14–15–18] with match time being the major determinant of this difference. However, comparisons across studies are difficult as researchers generally use generic descriptors for starters and nonstarters. For instance, assigning players into two or three accumulative load groups based on match time [14–15–18–28]. To complicate matters, studies use distinct loading metrics, thresholds, GPS devices, and these further limits comparability between studies. The present analysis is the most granular to date, and not only compares four groups of players based on match time but also explores this as a function of their playing position.

The data demonstrated that the 0-min and <60-min groups were lower for most external loading variables in comparison to >60-min and 90-min groups, with very little difference between the 90-min and the >60-min group. This finding could potentially justify the practices employed by some elite clubs that split the squad for MD+1 work based on greater or less than <60-min played in previous official matches [11]. For instance, professional Spanish team use a MD+1R session for players completing >60 min to aid recovery, as opposed to MD+1C that is for players completing <60 min and is a compensatory session in an attempt to replicate loading during an official match [11]. This is imperative, as matches seems to act as an important physiological, mechanical and neuromuscular stimulus for various fitness parameters [15–16–17]. As the MD+1C session encompassed both positional games and SSG in reduced pitch dimensions, it seems that this adequately mimicked the ACC and DEC match demands as differences were only present between the 90 and 0-min group [29]. This is not surprising as Martin-Garcia *et al.* [11] reported that ACC and DEC demands during a MD+1C session exceeded 80% of that performed in official match. Specifically, the ACC and DEC demands attained by the 90-min group equate to the load accumulated in 3.5 matches, which aligns with those reported by Stevens *et al.* [14], with the starter group also reaching more than three matches in high intensity ACC and DEC. In contrast, weekly ACC and DEC load in the 0-min group was equivalent to about three weekly matches, while other external load variables demonstrate weekly demands of between 1–2 matches. In agreement with the present findings, Stevens *et al.* [14] also found that ACC and DEC load variables displayed the lowest differential between starters and nonstarters compared to other external load variables.

Match running performances studies reveal that HSR and SPR are important physical actions, determinants many times for its connection with scoring opportunities [30], and also has been evidenced when comparing differences between performance level and playing positions [20–31–32]. The data demonstrate that HSR and SPR weekly accumulation was double for the 90-min vs the 0-min group. Despite this, the 0-min group still reached an average weekly accumulation value of one match for HSR and SPR. Interestingly, Stevens *et al.* [14] reported values of 1.5 and 0.5 for HSR and SPR for substitutes in one and two matches per week, respectively. Unfortunately, comparisons are difficult as this study only quantified one group for non-starters with no differentiation between substitutes and players who did not play at all. As previously discussed, strategies to replicate the match demands of nonstarters in MD+1C seems to be optimal for ACC and DEC [11]. However, the present results reveal that this session does not seem to provide sufficient overload to develop HSR and SPR capabilities as these external load variables only reach 20–30% of match play demands [11]. Research has indicated that pitch running without the ball or position specific circuits that encompass supra-maximal running alongside technical actions and some directional changes could achieve the necessary

overload for HSR and SPR metrics [33]. Although the relationship between the workload and the likelihood of injury is highly complex, influenced by different moderators [34] and in need of more research before conclusions can be drawn, weekly HSR and SPR management could be important [35]. This is especially important as a relationship between weekly HSR load and the probability of injury seems to be U-shaped whereby a moderate amount seems to have a protective effect [36]. Thus, perhaps both groups might present a certain risk of injury, in some cases due to under-stimulation (0-min group) and in other cases due to over-stimulation (90min group). Thus, future research should modify MD+1C sessions using a multi-modal approach to optimise the training stimulus for players with partial or no limited match time.

The tactical role of a player seems to be a powerful determinant of their match physical performance, so it is imperative that the conditioning stimulus has a positional element to it [2]. It is worth noting that during official matches of this same team studied, CB and MF (~300–400 m) cover less HSR than FW and FB (~600–800 m). Interestingly, FB produce the greatest HSR and SPR load and MF typically the lowest load compared to other positions from MD-4 across to MD-1 [11]. Thus, training load and match running metrics are directly aligned to enable players to be conditioned for their distinct tactical role in the team. Although interpreting positional trends from a low sample size requires caution, it was clear to see that this trend was not always evident for various metrics across the various weekly accumulative load groups. Crucially, as football is a submaximal sport whereby players are likely to be working within their physical capacity for long periods, it is imperative to remember that players tax a high proportion of this capacity during intense periods of the match [37]. Thus, inadequate match time could impact a player's ability to maintain a high work rate if required during various phases of match-play. As selected positions within the modern game have higher physical requirements [19], it might be wise to ensure these positions are loaded more during compensatory sessions when not gaining match time. This is especially important for FB to enable them to cope with a dual tactical role that includes defensive (e.g. recovery running) and offensive (e.g. overlapping and running the channel) physical-tactical actions [2]. Research also highlights that most substitutions occurred at halftime and between the 60- to 85-min [38] and these include more attacking positions as the second half progresses [21]. Thus, as offensive players are more likely to gain some match time, bespoke conditioning work might need to be incorporated to account for defensive players not been regularly introduced and attaining 0 min per week regularly.

Some of the main limitations of the study are due to not considering the specific load in the training made outside the field (indoor), not incorporating internal load variables (heart rate or RPE) or the effective differences in minutes that may exist within the same group (61 minutes vs. 89 minutes). However, the use of four groups depending on the minutes played in the previous match (90 minutes, >60 minutes, <60 minutes and 0 minutes) provides new and practical information to the coaches.

## CONCLUSIONS

In summary, this study revealed that players with more match playing time had a higher accumulative weekly load for most of the variables, but particularly at HSR and SPR. Moreover, position specific accumulative load trends were evident in players completing more match time, with less positional variation when expressed relative to match requirements.

## Practical applications

Knowing external training loads is important for applied professionals, particularly when it comes to optimizing position-specific loads. As competitive match play is an important stimulus for developing the physiological capacities of players regularly completing games, it is imperative that practical strategies are implemented to offset any reductions in the fitness of players getting limited game time. Thus, the MD-1, MD and MD+1 sessions could be the most appropriate sessions to try to compensate for the absence of competitive stimulus in order to synchronize the weekly training loads of all the team members with a view to having all the effective with the fitness and the necessary freshness with which to be able to compete every week.

## Acknowledgments

The authors thank all the athletes who participated in the study. The authors did not report any potential conflicts of interest. The authors gratefully acknowledge the support of a Spanish government subproject Mixed method approach on performance analysis (in training and competition) in elite and academy sport [PGC2018-098742-B-C33] (2019-2021) [del Ministerio de Ciencia, Innovación y Universidades (MCIU), la Agencia Estatal de Investigación (AEI) y el Fondo Europeo de Desarrollo Regional (FEDER)], that is part of the coordinated project New approach of research in physical activity and sport from mixed methods perspective (NARPAS\_MM) [SPGC201800X098742CV0].

## Conflict of interest

The authors did not report any potential conflicts of interest.

## REFERENCES

1. Bradley PS, Ade JD. Are current physical match performance metrics in elite soccer fit for purpose or is the adoption of an integrated approach needed? *Int J Sports Physiol Perform.* 2018; 13(5):656–664.
2. Ade J, Fitzpatrick J, and Bradley PS. High-intensity efforts in elite soccer matches and associated movement patterns, technical skills and tactical actions. *Information for position-specific training drills.* *J Sports Sci.* 2016; 34(24):2205–2214.
3. Bradley PS, Mascio MD, Mohr M, Fransson D, Wells C, Moreira A, et al. Can modern trends in elite football match demands be translated into novel training

- and testing modes? *Aspartar Sports Med J.* 2018;7(1):46–52.
4. Moreira A, Bradley P, Carling C, Arruda AF, Spigolon LM, Franciscan C, Aoki MS. Effect of a congested match schedule on immune-endocrine responses, technical performance and session-RPE in elite youth soccer players. *J Sports Sci.* 2016;34(24): 2255–2261.
  5. Carling C, Le Gall F, McCall A, Nedelec M, Dupont G. Squad management, injury and match performance in a professional soccer team over a championship-winning season. *Eur J Sport Sci.* 2015; 15(7):573–582.
  6. Gabbett T. The training-injury prevention paradox: Should athletes be training smarter and harder? *Br J Sports Med.* 2016;50(5):273–280.
  7. Drew MK, Finch CF. The relationship between training load and injury, illness and soreness: a systematic and literature review. *Sports Med.* 2016; 4(6):861–883.
  8. Gabbett TJ, Whiteley R. Two Training-Load Paradoxes: Can We Work Harder and Smarter, Can Physical Preparation and Medical Be Teammates? *Int J Sports Physiol Perform.* 2017;12(s2):S2–50.
  9. Coutts A, Reaburn P, Piva TJ, Murphy A. Changes in selected biochemical, muscular strength, power, and endurance measures during deliberate overreaching and tapering in rugby league players. *Int J Sports Med.* 2007;28(02):116–124.
  10. Fessi MS, Zarrouk N, Di Salvo V, Filetti C, Barker AR, Moalla W. Effects of tapering on physical match activities in professional soccer players. *J Sport Sci.* 2016;34(24):2189–2194.
  11. Martín-García A, Gómez Díaz A, Bradley PS, Cos F, Casamichana D. The Quantification of a Professional Football Teams External Load Using a Microcycle Structure. *J Strength Cond Res.* 2018; 32(12):3511–3518.
  12. Owen AL, Djaoui L, Newton M, Malone S, Mendes B. A contemporary multi-modal mechanical approach to training monitoring in elite professional soccer. *Sci Med Football.* 2017; 1(3):216–221.
  13. Akenhead R, Harley JA, Tweddle SP. Examining the external training load of an English Premier League football team with special reference to acceleration. *J Strength Cond Res.* 2016; 30(9):2424–2432.
  14. Stevens TG, de Ruiter CJ, Twisk JW, Savelsbergh GJ, Beek PJ. Quantification of in-season training load relative to match load in professional Dutch Eredivisie football players. *Sci Med Football.* 2017;1(2):117–125.
  15. Anderson L, Orme P, Di Michele R, Close GL, Morgans R, Drust B, et al. Quantification of training load during one-, two-and threeweek schedules in professional soccer players from the English Premier League: Implications for carbohydrate periodisation. *J Sport Sci.* 2016;34(13):1250–1259.
  16. Morgans R, Di Michele R, Drust B. Soccer match play as an important component of the power-training stimulus in Premier League players. *Int J Sports Physiol Perform.* 2017;13(5):665–667.
  17. Young WB, Newton RU, Doyle TLA, Chapman D, Cormack S, Stewart C, et al. Physiological and anthropometric characteristics of starters and non-starters and playing positions in elite Australian Rules Football: A case study. *J Sci Med Sport.* 2005; 8(3):333–345.
  18. Azcárate U, Yanci J, Los Arcos A. Influence of match playing time and the length of the between-match microcycle in Spanish professional soccer players' perceived training load. *Sci Med Football.* 2018;2(1):23–28.
  19. Bush, MD, Archer, DT, Hogg, R, and Bradley, PS. Factors influencing physical and technical variability in the English Premier League. *Int J Sports Physiol Perform.* 2015;10(7):865–872.
  20. Bradley PS, Sheldon W, Wooster B, Olsen P, Boanas P, Krustup P. High-intensity running in English FA Premier League soccer matches. *J Sport Sci.* 2009;27(2):159–168.
  21. Bradley PS, Lago-Peñas C, Rey E. Evaluation of the match performances of substitution players in elite soccer. *Int J Sports Physiol Perform.* 2014; 9(3):415–424.
  22. Lacombe M, Simpson BM, Cholley Y, Lambert P, Buchheit M. Small-sided games in elite soccer: Does one size fits all? *Int J Sports Physiol Perform.* 2018; 13(5):568–576.
  23. Malone JJ, Di Michele R, Morgans R, Burgess D, Morton JP, Drust B. Seasonal training-load quantification in elite English premier league soccer players. *Int J Sports Physiol Perform.* 2015; 10(4):489–497.
  24. Beato M, Bartolini D, Ghia G, Zamparo P. Accuracy of a 10 Hz GPS unit in measuring shuttle velocity performed at different speeds and distances (5–20 M). *J Hum Kinet.* 2016;54(1):15–22.
  25. Buchheit M, Simpson BM. Player-tracking technology: Half-full or half-empty glass? *Int J Sports Physiol Perform.* 2017;12(s2):S2–35.
  26. Castellano J, Alvarez-Pastor D, Bradley PS. Evaluation of research using computerised tracking systems (Amisco and Prozone) to analyse physical performance in elite soccer: A systematic review. *Sports Med.* 2014; 44(5):701–712.
  27. Batterham AM, Hopkins WG. Making meaningful inferences about magnitudes. *Int J Sports Physiol Perform.* 2006;1(1):50–57.
  28. Los Arcos A, Mendez-Villanueva A, Martínez-Santos R. In-season training periodization of professional soccer players. *Biol Sport.* 2017; 34(2):149–155.
  29. Castellano J, Casamichana D. Differences in the number of accelerations between small-sided games and friendly matches in soccer. *J Sports Sci Med.* 2013; 12(1):209–210.
  30. Hinijsa A, Castellano J. Influencia de la distancia recorrida en diferentes rangos de velocidad en la anotación de un gol en fútbol. *Retos.* 2017;31(1):188–192.
  31. Buchheit M, Simpson BM, Hader K, Lacombe M. Occurrences of near-to-maximal speed running bouts in elite soccer: insights for training prescription and injury mitigation. *Sci Med.* 2020; In press.
  32. Mohr M, Krustup P, Bangsbo J. Match performance of high- standard soccer players with special reference to development of fatigue. *J Sports Sci.* 2003;21(7):519–528.
  33. Ade J, Harley J, Bradley PS. The physiological response, time-motion characteristics and reproducibility of various speed endurance drills in elite youth soccer players: Small sided games vs generic running. *Int J Sports Physiol Perform.* 2014;9(3):471–479.
  34. Windt J, Zumbo BD, Sporer B, MacDonald K, Gabbett TJ. Why do workload spikes cause injuries, and which athletes are at higher risk? Mediators and moderators in workload–injury investigations. *Br J Sports Med.* 2017;993–994.
  35. Hader K, Rumpf MC, Hertzog M, Kilduff LP, Girard O, Silva JR. Monitoring the Athlete Match Response: Can External Load Variables Predict Post-match Acute and Residual Fatigue in Soccer? A Systematic Review with Meta-analysis. *J Sports Med.* 2019; 5(1):48.
  36. Malone S, Owen A, Mendes B, Hughes B, Collins K, Gabbett T. High-speed running and sprinting as an injury risk factor in soccer: Can well-developed physical qualities reduce the risk? *J Sci Med Sport.* 2017;21(3):257–262.
  37. Paul DJ, Bradley PS, Nasis G. Factor affecting running performance in elite soccer players: Shedding some light on the complexity. *Int J Sports Physiol Perform.* 2015;10(4):516–519.
  38. Rey E, Lago-Ballesteros J, Padrón-Cabo A. Timing and tactical analysis of player substitutions in the UEFA Champions League. *Int J Perf Anal Spor.* 2015;15(3):840–850.